



## Nature of O<sub>2</sub> and CO Binding to Metalloporphyrins and Heme Proteins

James P. Collman; John I. Brauman; Thomas R. Halbert; Kenneth S. Suslick

*Proceedings of the National Academy of Sciences of the United States of America*,  
Volume 73, Issue 10 (Oct., 1976), 3333-3337.

Stable URL:

<http://links.jstor.org/sici?sici=0027-8424%28197610%2973%3A10%3C3333%3ANOOACB%3E2.0.CO%3B2-2>

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

*Proceedings of the National Academy of Sciences of the United States of America* is published by National Academy of Sciences. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/nas.html>.

---

*Proceedings of the National Academy of Sciences of the United States of America*  
©1976 National Academy of Sciences

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact [jstor-info@umich.edu](mailto:jstor-info@umich.edu).

©2003 JSTOR

<http://www.jstor.org/>  
Thu Apr 3 18:10:51 2003

# Nature of O<sub>2</sub> and CO binding to metalloporphyrins and heme proteins

(hemoglobin/infrared vibrational spectra/metal bound superoxides/distal binding site)

JAMES P. COLLMAN, JOHN I. BRAUMAN, THOMAS R. HALBERT, AND KENNETH S. SUSLICK

Department of Chemistry, Stanford University, Stanford, California 94305

Contributed by James P. Collman, July 12, 1976

**ABSTRACT** The O<sub>2</sub> vibration of dioxygen adducts of Fe and Co model complexes of  $\alpha,\alpha,\alpha,\alpha$ -tetrapivalamidophenylporphyrin ("picket fence" porphyrin, TpivPP) with 1-methylimidazole and 1-tritylimidazole as axial bases are reported, obtained with difference techniques between <sup>16</sup>O<sub>2</sub>, <sup>18</sup>O<sub>2</sub>, <sup>16</sup>O-<sup>18</sup>O, and NO with a Fourier transform infrared spectrometer. Assignments of  $\nu_{O_2}$  are (O<sub>2</sub>)Fe(TpivPP) 1-methylimidazole, 1159 cm<sup>-1</sup> in Nujol; (O<sub>2</sub>)Fe(TpivPP) 1-tritylimidazole, 1163 in benzene; (O<sub>2</sub>)Co(TpivPP) 1-methylimidazole, 1150 in Nujol; (O<sub>2</sub>)Co(TpivPP) 1-tritylimidazole, 1153 in benzene. Comparisons with other known Fe, Co, Cr, and Ti dioxygen complexes are made, and it is concluded that the bent dioxygen ligand is best viewed as bound superoxide, O<sub>2</sub><sup>-</sup>.

The CO affinities of various hemoproteins and model systems are discussed. A correlation between the CO stretching frequency and its binding constant is described. The drastically lowered affinity of hemoproteins for CO compared with unencumbered models is attributed to steric hindrance in the distal binding site, which allows discrimination between the already bent Fe<sup>III</sup>-O<sub>2</sub><sup>-</sup> and the normally linear Fe<sup>II</sup>-CO systems. If the affinity of hemoproteins in living systems for CO relative to O<sub>2</sub> were not decreased, then massive poisoning would result from endogenous CO.

A continuing controversy exists over the nature of ligand bonding to hemoproteins and metalloporphyrins. While it has been well established that both iron and cobalt bind dioxygen in a bent end-on fashion (1, 2), physical measurements (3-6) and theoretical calculations (7-10) have been used to support both the spin-paired dioxygen (M<sup>II</sup>-O<sub>2</sub>) (11) and the electron transfer superoxide (M<sup>III</sup>-O<sub>2</sub><sup>-</sup>) (12) models. Here we present infrared data on dioxygen "picket-fence" porphyrin complexes of iron and cobalt. These results strongly support the M<sup>III</sup>-O<sub>2</sub><sup>-</sup> formalism. The present data correct the preliminary report of an oxygen vibration at 1385 cm<sup>-1</sup> (3, 13) which has since proved to be an artifact\* (14). We are unaware of any previous reports of  $\nu_{O_2}$  in model Fe and Co porphyrins. In addition, we discuss the nature of CO binding to ferrous porphyrins and the implication this has for the role of the apoprotein in biological oxygen carriers. It is our thesis that the distal binding site in hemoproteins functions to lower the carbon monoxide affinity.

## MATERIALS AND METHODS

Benzene (spectroquality, Baker "Photrex") solutions of the "picket fence" porphyrin complexes (3, 14), Fe<sup>II</sup>meso-tetra( $\alpha,\alpha,\alpha,\alpha$ -pivalamidophenyl)porphyrinato (TpivPP) and Co<sup>II</sup> ( $\alpha,\alpha,\alpha,\alpha$ -TpivPP), 17 mM and 19 mM containing 5 and 2 equivalents of 1-tritylimidazole (NTrIm), respectively, were

Abbreviations: TpivPP, meso-tetra( $\alpha,\alpha,\alpha,\alpha$ -pivalamidophenyl)porphyrinato; NTrIm, 1-tritylimidazole; NMeIm, 1-methylimidazole; P<sub>1/2</sub>, partial pressure of gas at half-saturation; py, pyridine; one torr = 1.33 × 10<sup>2</sup> pascal.

\* This band at 1385 cm<sup>-1</sup> is due to an impurity found in occasional KBr samples (R. R. Gagne, personal communication).

prepared under inert atmosphere. Separate samples of these solutions were exposed to an atmosphere of CO, NO, O<sub>2</sub>, and >200 torr <sup>18</sup>O<sub>2</sub> (both 54% and 90% enriched)<sup>†</sup>. Infrared spectral data of all solutions were collected between 2000 cm<sup>-1</sup> and 750 cm<sup>-1</sup> on a Digilab FTS 15 B Fourier transform spectrometer, at 1 cm<sup>-1</sup> resolution. A RRIC XL0 variable pathlength cell fitted with NaCl windows and set to 0.1 mm was used. Difference spectra were computed using the standard software of the spectrometer. (These difference spectra are analogous to sample minus reference absorbance spectra on a conventional dual beam spectrometer.) The coefficients in the difference spectra were chosen to reduce to baseline the porphyrin bands at 1695 cm<sup>-1</sup> and 1001 cm<sup>-1</sup>; small amounts of a neat benzene spectrum were then added or subtracted, to account for slight changes in concentration and pathlength, so as to reduce to baseline the benzene bands at 1961 cm<sup>-1</sup> and 1819 cm<sup>-1</sup>. In addition, solid samples of Fe(TpivPP)(NMeIm) and Co(TpivPP)(NMeIm) exposed to O<sub>2</sub>, <sup>18</sup>O<sub>2</sub> (90% enriched), and NO were prepared as Nujol mulls between KBr plates; difference spectra were computed as above at 4 cm<sup>-1</sup> resolution.

## RESULTS

The spectrum of Fe(TpivPP)(NTrIm)O<sub>2</sub> in benzene versus benzene is shown in Fig. 1. The gaps left at 1480-1460 cm<sup>-1</sup> and 1040-1020 cm<sup>-1</sup> are due to the most intense bands of benzene whose absorbances in these regions exceed 99.9% under these conditions. The spectrum in benzene, apart from increased resolution, does not differ from those taken in KBr pellets and Nujol mulls.

Difference spectra of the Fe(TpivPP)(NTrIm) adducts of <sup>16</sup>O<sub>2</sub> versus <sup>18</sup>O<sub>2</sub> (both 90% and 54% enriched) are shown in Fig. 2. Similar spectra for the Co analog are shown in Fig. 3. The bands due to <sup>16</sup>O<sub>2</sub>, <sup>16</sup>O-<sup>18</sup>O, and <sup>18</sup>O<sub>2</sub> are evident, as is an unassigned feature in the region of 1110-1095 cm<sup>-1</sup>. In the O<sub>2</sub> adduct versus NO adduct spectra, one observes the complete loss of the 1163 cm<sup>-1</sup> peak in Fe and the 1153 cm<sup>-1</sup> peak in Co and the appearance of new bands at 1080 cm<sup>-1</sup> and 1077 cm<sup>-1</sup>, respectively, upon going to <sup>18</sup>O<sub>2</sub> (90%) which confirms the assignment of these bands. The shifts in stretching frequencies of <sup>16</sup>O<sub>2</sub> and <sup>16</sup>O-<sup>18</sup>O adducts of both Fe and Co match the harmonic oscillator prediction. However, the band due to <sup>18</sup>O<sub>2</sub> is shifted to a considerably lower frequency than predicted. This, in conjunction with the unassigned feature in the 1100 cm<sup>-1</sup> region, both in benzene solution and in Nujol mull, indicates that Fermi coupling, i.e., the splitting of an accidental

<sup>†</sup> Recent studies in this laboratory on the binding constants of O<sub>2</sub> with both Fe(TpivPP) 1-methylimidazole (NMeIm) and Co(TpivPP) (NMeIm) show that the partial pressure of O<sub>2</sub> at half saturation (P<sub>1/2</sub>O<sub>2</sub>) = 0.31 torr and P<sub>1/2</sub>O<sub>2</sub> about 100 torr, respectively, at room temperature. While both Fe and Co complexes bind NO, only Fe binds CO as a six coordinate complex.




FIG. 1.  $\text{Fe}(\text{TpivPP})(\text{NTrIm})^{16}\text{O}_2$  full spectra in benzene versus benzene.

degeneracy (15–18) has occurred between the strong porphyrin bands in this region and some of the isotopic  $\text{O}_2$  stretches. The intensity of the observed  $^{16}\text{O}_2$  bands decreases in the expected amount (to within experimental error) between the  $^{16}\text{O}_2$  versus  $^{18}\text{O}_2$  (90%) and the  $^{16}\text{O}_2$  versus  $^{18}\text{O}_2$  (54%) spectra. A more detailed analysis of these spectra is not possible at this time; however, the main points are clear. The  $\text{O}_2$  stretch in both the Fe and the Co “picket-fence” porphyrin complexes is in the region of  $1150\text{ cm}^{-1}$ , is relatively independent of the metal, and has been observed both in benzene solution and in Nujol mull.

Table 1 summarizes the observed band positions of the  $\text{O}_2$  stretch, together with data from several other relevant systems. Table 2 presents the stretching frequencies and binding constants of the CO adducts of biological and model Fe porphyrins.

## DISCUSSION

**Nature of  $\text{O}_2$  Binding.** The frequencies of the  $\text{O}_2$  vibration in our model porphyrins are close to those of ionic superoxides, whose range is about  $1150\text{--}1100\text{ cm}^{-1}$  (29), with a gap of  $200\text{ cm}^{-1}$  from the other dioxygen species,  $\text{O}_2$  and  $\text{O}_2^-$ . The only




FIG. 2. Difference spectra of  $\text{Fe}(\text{TpivPP})(\text{NTrIm})^{16}\text{O}_2$  minus  $\text{Fe}(\text{TpivPP})(\text{NTrIm})^{18}\text{O}_2$ . Spectrum *a* taken with 90% enriched  $^{18}\text{O}_2$ ; spectrum *b* taken with 54% enriched  $^{18}\text{O}_2$  in benzene solution. Extinction coefficient of  $^{16}\text{O}_2$  band is  $320\text{ M}^{-1}\text{ cm}^{-1}$  with half height width of  $10\text{ cm}^{-1}$  and an integrated intensity  $B = 3700\text{ M}^{-1}\text{ cm}^{-2}$ . For comparison, the CO vibration in  $\text{Fe}(\text{TpivPP})(\text{NTrIm})\text{CO}$  has  $\epsilon = 730\text{ M}^{-1}\text{ cm}^{-1}$  and a half height width of  $10\text{ cm}^{-1}$  and an integrated intensity  $B = 7500\text{ M}^{-1}\text{ cm}^{-2}$ . Estimated errors in the intensity data are less than 20%. Gap left at  $1020\text{--}1040\text{ cm}^{-1}$  due to intense benzene absorption.




FIG. 3. Difference spectra of  $\text{Co}(\text{TpivPP})(\text{NTrIm})^{16}\text{O}_2$  minus  $\text{Co}(\text{FeTpivPP})(\text{NTrIm})^{18}\text{O}_2$ . Spectrum *a* taken with 90% enriched  $^{18}\text{O}_2$ ; spectrum *b* taken with 54% enriched  $^{18}\text{O}_2$  in benzene solution. The bands attributed to the  $\text{O}_2$  vibration are doublets resolved by  $5\text{ cm}^{-1}$ :  $1150$  and  $1155\text{ cm}^{-1}$  for  $^{16}\text{O}_2$  and  $1179$  and  $1174\text{ cm}^{-1}$  for  $^{18}\text{O}_2$ . Gap left at  $1020\text{--}1040\text{ cm}^{-1}$  due to intense benzene absorption.

Table 1. Vibrational frequencies of dioxygen complexes

Compound	$\nu_{O_2}$ (cm <sup>-1</sup> )	Ref.
Fe (TpivPP)		
(NTrIm)O <sub>2</sub> *	1163 (1080, 1127 <sup>†</sup> )	
Fe (TpivPP)		
(NMeIm)O <sub>2</sub> ‡	1159 (1075)	
MbO <sub>2</sub>	1103 (1065)	18
HbO <sub>2</sub>	1107 (1065)	19
Hb(deutFe)O <sub>2</sub>	1106 (1065)	20
Co(TpivPP)		
(NTrIm)O <sub>2</sub> *	1153§ (1077§, 1125)	
Co(TpivPP)		
(NMeIm)O <sub>2</sub> ‡	1150 (1077)	
Hb(deut Co)O <sub>2</sub>	1105 (1065)	20
Co(acen)(py)O <sub>2</sub>	~1130	21
Co(bzacen)(py)O <sub>2</sub>	1128	22
[Co(CN) <sub>5</sub> O <sub>2</sub> ] <sup>-3</sup>	1138	23
Cr(TPP)(py)O <sub>2</sub>	1142	24
Ti(OEP)O <sub>2</sub>	898	25
ML <sub>2</sub> (CO)(X)O <sub>2</sub> ¶	830–900	26
O <sub>2</sub> <sup>+</sup>	1860	17
O <sub>2</sub>	1556	17, 27
O <sub>2</sub> ( <sup>1</sup> Δ)	1484	17
O <sub>2</sub> <sup>-</sup>	1150–1100	17, 28, 29
O <sub>2</sub> <sup>=</sup>	740–850	17, 27, 29

Acen, *N,N'*-ethylenebis(acetylacetoniminato); [bzacen, benzyl-*N,N'*-ethylenebis(acetylacetoniminato)]; py, pyridine; TPP, *meso*-tetraphenylporphyrinato; OEP, octaethylporphyrinato; deut, deuteroporphyrinato.

\* Benzene solution.

† Wavenumber for <sup>18</sup>O<sub>2</sub> and <sup>16</sup>O–<sup>18</sup>O, respectively.

‡ Nujol mull.

§ These are, in fact, doublets with  $\nu(^{16}O_2) = 1155, 1150$  cm<sup>-1</sup> and  $\nu(^{18}O_2) = 1179, 1174$  cm<sup>-1</sup> due perhaps to the two different conformers of O<sub>2</sub> and NMeIm orientations.

¶ For example, M = Ir, Co, Os, Rh, Ni, Pt, Pd; L = triphenylphosphine, triphenylarsine, dialkylphenylphosphine; X = none, Cl, Br, I, N<sub>3</sub>, NO<sub>3</sub>.

other iron bound O<sub>2</sub> systems with which to compare are Hb and Mb; as we see in Table 1, all are in the spectral region associated with superoxide. However, the 50 cm<sup>-1</sup> difference in the dioxygen vibration of the hemoproteins and the model [Fe(TpivPP)(NMeIm)O<sub>2</sub>] is larger than would be expected on the basis of *electronic* differences between a tetraaryl and a natural porphyrin. For example,  $\nu_{NO}$  for Fe(dimethyl protoporphyrinato)(NMeIm)NO is 1630 cm<sup>-1</sup> in CHCl<sub>3</sub> or CH<sub>2</sub>ClCH<sub>2</sub>Cl (39), while for Fe(*meso*-tetra(pivalamidophenyl)porphyrinato)(NMeIm)NO is 1632 cm<sup>-1</sup> in benzene or Nujol mull. In addition,  $\nu_{CO}$  for Fe(dimethyl protoporphyrinato IX)(NMeIm)CO is 1970 cm<sup>-1</sup> in KBr (37) and 1969 cm<sup>-1</sup> in CHCl<sub>3</sub> (30, 39) and for Fe(TpivPP)(NMeIm)CO 1967 ± 2 cm<sup>-1</sup> in benzene, Nujol, KBr, or CH<sub>2</sub>Cl<sub>2</sub>. One would expect, then, that the frequencies of the O<sub>2</sub> vibration would not change significantly in going from TpivPP to protoporphyrinato (i.e., the heme of Hb and Mb) *in the absence of* other differences in the local environment of the binding site such as steric interactions and local polarity. Because the actual normal mode which contains the O<sub>2</sub> stretch includes small amounts of other bends and stretches, small changes in geometry, as well as polarity, will cause shifts in the observed frequency, and the shift between the model complex and the hemoproteins, while interesting, is not surprising. The 1150 cm<sup>-1</sup> band of the Co "picket fence" porphyrin complex is entirely consistent with the frequencies found for nonpor-

Table 2. Carbon monoxide adducts of Fe porphyrins

Compound	$\nu_{CO}$ (cm <sup>-1</sup> )	P <sub>1/2</sub> CO*, 20°–25°, pH 7 (torr)	Ref.
MbCO <sup>†</sup>	1945	0.018	30, 31
HbCO <sup>†</sup>	1951	0.004 (0.035)‡	30, 31, 32, 33
HbCO, $\alpha$ or $\beta$ chains		0.002	31
HbCO, insect (chironimus)	1959	0.002	34, 35
HbCO Zurich ( $\beta_{63}$ His → Arg)	1951, 1958	P <sub>1/2</sub> $\beta$ < 1/3 P <sub>1/2</sub> $\alpha$	36
Fe(P) (NMeIm) CO	1970§	0.0002¶	37, 38
Fe(TpivPP) (NMeIm)CO	1969	"Irreversible"¶¶	

\* Equilibrium constants involving dissolved gases should be expressed in terms of the partial pressure of the gas over the solution, not the concentration of the gas in solution. Because the activity of the dissolved gas is equal to the partial pressure of the gas above the solution, direct comparisons can be made of K<sub>eq</sub> in various solvents, without regard to gas solubility, *only if* K<sub>eq</sub> is expressed in terms of pressure.

† These values are typical for a wide range of Mbs and Hbs.

‡ These are the P<sub>1/2</sub>CO for the binding of the fourth CO and for the overall equilibrium, respectively.

§ P = diethylprotoporphyrinato.

¶ P = deuteroporphyrinato in benzene.

¶¶ Evacuation at 10<sup>-5</sup> torr for 75 hr does not affect the intensity of the  $\nu_{CO}$  bond. Hb Zurich refers to hemoglobin Zurich.

phyrin mononuclear Co–O<sub>2</sub> adducts, as listed in Table 1. The small shift (10 cm<sup>-1</sup>) between the  $\nu_{O_2}$  in our Fe and Co complexes lends strong support to Maxwell and Caughey's surprising report of a shift of only 1 cm<sup>-1</sup> between deuteroporphyrin Fe and Co reconstituted oxy-Hb.

In summary, the O<sub>2</sub> vibrational frequencies in our complexes are in the superoxide region. The influence of the metal is minimal. In conjunction with the electron spin resonance (6), x-ray photoelectron spectroscopy (5), and bond lengths from x-ray diffraction data (2, 41, 42), this and prior infrared evidence convincingly demonstrate that, for Co, 1:1 dioxygen adducts are best formulated as Co<sup>III</sup>-O<sub>2</sub><sup>-</sup>. It is now difficult to escape a similar conclusion for Fe and Cr (24). Thus, the best current picture is that the bent dioxygen ligand should be viewed as bound superoxide.

**Nature of CO Binding.** The infrared stretching frequencies of bound CO in hemes and hemoproteins appear to correlate with the equilibrium constants for binding of CO. More importantly, the hemoprotein structure functions to *reduce* significantly the affinity for CO. Table 2 presents the binding constants and stretching frequencies for a series of Fe porphyrin CO adducts. It has recently been suggested (42) that the  $\nu_{CO}$  of ferrous porphyrin proteins correlates with the redox potential for the Fe<sup>III</sup>/Fe<sup>II</sup> couple. However, this correlation fails to hold in simple porphyrin systems: as noted above,  $\nu_{CO}$  is not affected by the nature of the porphyrin, whereas E<sub>1/2</sub>Fe<sup>III</sup>/Fe<sup>II</sup> changes by more than 100 mV (43) where E<sub>1/2</sub> refers to the electrochemical half-wave potential. We suggest that the lowering of  $\nu_{CO}$  in Mb and Hb from that in simple porphyrins is, at least in part, due to distortion of the linear Fe–CO group from a normal to the porphyrin plane by bulky amino acid residues near the binding site. This tilted conformation is now documented by structure determinations on three diverse hemoproteins: insect (distal residue: Ile) (44), bloodworm (distal residues: Leu, Val)

(45), and sperm whale Mb (distal residues: His, Val) (46). In simple porphyrins the CO should be linear, as in the structure of Fe(TpivPP)(NMeIm)CO (47).

The hypothesis that the  $\nu_{\text{CO}}$  is altered by distortion is supported by several pieces of evidence which indicate that  $\nu_{\text{CO}}$  can be altered by affecting the structure of the binding site of hemoproteins. Denatured Hb in a KBr pellet has  $\nu_{\text{CO}} = 1970 \text{ cm}^{-1}$  (37), the same as unhindered porphyrins. Mutant Hbs have  $\nu_{\text{CO}}$  approaching those of free porphyrins if the distal histidine is replaced (48): HbZ ( $\beta_{63} \text{ His} \rightarrow \text{Arg}$ ) shows  $\nu_{\text{CO}}$  at  $1951 \text{ cm}^{-1}$  (the normal  $\alpha$  chains) and  $1958 \text{ cm}^{-1}$  (the  $\beta$  chains) and HbM<sub>Emory</sub> ( $\beta_{63} \text{ His} \rightarrow \text{Tyr}$ ) at  $1951 \text{ cm}^{-1}$  and  $1970 \text{ cm}^{-1}$ . In these mutants the binding site is thought to be more open (49).

Furthermore, the CO stretching frequency and the CO affinity constant appear to correlate. This correlation has been overlooked in the past (48) because the overall affinity of Mb for CO is higher than that of Hb, while the  $\nu_{\text{CO}}$  is reversed: however, because  $\nu_{\text{CO}}$  is probing the *relaxed* conformation of Hb, it is *not* the *overall* CO affinity which one should examine, but that for binding to the relaxed form. This can be approximated by the intrinsic binding constant (including the statistical factor of 4) for binding the fourth CO,  $\text{Hb}(\text{CO})_3 \rightleftharpoons \text{Hb}(\text{CO})_4$ , which is 10-fold greater in its affinity for CO than the overall Hb equilibrium constant and hence four times *greater* than that of Mb. In addition, the excellent work of Caughey and associates (36) indicates that hemoglobin Zurich binds CO at least three times stronger in its mutant  $\beta$  chains ( $\nu_{\text{CO}} = 1958 \text{ cm}^{-1}$ ) than in its normal  $\alpha$  chains ( $1951 \text{ cm}^{-1}$ ). Table 2 gives persuasive data that the lower the  $\nu_{\text{CO}}$ , the lower the affinity for CO. There are, of course, other significant contributions to  $\nu_{\text{CO}}$  (especially the local polarity). The portion which each of these play must, at the moment, remain in doubt.

It is our hypothesis that a major function of the protein is to provide steric bulk to the ligand binding site so as to distort normally linear ligand systems (e.g., Fe-CO). It must be stressed how remarkable it is that three separate hemoproteins with quite different amino acid sequences—even in the immediate vicinity of the binding site—all show distorted CO binding. The effect of this distortion is readily apparent in the CO affinity constants (compare  $P_{1/2}^{\text{CO}}$  in Table 2). Between the free porphyrin in benzene and Mb there is a 100-fold difference in the binding of  $\text{CO}^\ddagger$  while there is only a 2-fold difference in the binding of the intrinsically bent  $\text{O}_2^-$  (14). Other effects of the distal residues (especially histidine) such as increased polarity, hydrogen bonding, or other bonding interactions (53) serve only to stabilize the binding of CO.

The teleological rationale for the reduced CO affinity in hemoproteins stems from the high concentration of CO which biological systems must face. Although the partial pressure of CO,  $P_{\text{CO}}$ , present in the uncontaminated atmosphere [less than 0.2 ppm (54)] is insignificant even to unconstrained ferrous heme models, the primary source of CO in biological systems is *endogenous*: the use of porphyrins and the production of CO are inherently linked. The biological catabolism of Hb and Mb has been known for more than 20 years to produce 1 mol of CO per heme (55–57) with the production of biliverdin-IX $\alpha$  which undergoes further decomposition to bilirubin, the yellow bile pigment. It is important that this catabolic pathway appears

to be a kinetically and thermodynamically intrinsic property of iron porphyrins; even hemin, merely in the presence of ascorbic acid and oxygen, will produce a stoichiometric quantity of CO (57, 58). The catabolism of Hb, Mb, P-450, cytochromes, and other porphyrin containing proteins causes, even in air with  $P_{\text{CO}} = 0$ , roughly 1% of the body's Hb and Mb to be bound as HbCO and MbCO (59). In a closed system, the equilibrium  $P_{\text{CO}}$  in mammalian tissue is 10–50 ppm (i.e., at this ambient  $P_{\text{CO}}$  animals show no net loss or gain in  $[\text{HbCO}]$ ), corresponding to 4–5% COHb (56). The tilting of CO in biological oxygen carriers such as Hb and Mb assumes critical importance in light of the inherent CO production from porphyrin catabolism. If Hb and Mb did not have hindered binding sites for CO and if CO were not forced to distort, then the amount of Hb and Mb poisoned by endogenous CO would be considerably greater. For unhindered Fe porphyrins,  $[\text{FeCO}]/[\text{FeO}_2](P_{\text{O}_2}/P_{\text{CO}})$ , commonly referred to as  $M$ , is greater than 1500 (from Table 2). With a local  $P_{\text{CO}}$  about 30 ppm,  $[\text{HbCO}]$  or  $[\text{MbCO}]$  would be 20% in the absence of steric hindrance, at which level heavy work becomes difficult and functioning is impaired. This lowered CO affinity of hemoproteins is, of course, independent of any possible correlation with  $\nu_{\text{CO}}$ .

Several experiments are suggested by these observations. Affinity constants for CO for the mutant hemoglobins Hb Zurich and HbM<sub>Emory</sub> (=HbM<sub>Saskatoon</sub>) or HbM<sub>Boston</sub> ( $\alpha_{58} \text{ His} \rightarrow \text{Tyr}$ ), which apparently have not been measured, would clarify the potential correlation of  $\nu_{\text{CO}}$  with the affinity constant [although the on-rates of recombination have been measured (60, 61) and are much faster than HbA, as expected]. The difficult experiment of measuring  $\nu_{\text{O}_2}$  in such mutant Hbs whose distal pocket is more open than in HbA would shed light on the source of the  $50 \text{ cm}^{-1}$  difference between our model dioxygen adducts'  $\nu_{\text{O}_2}$  and that of Hb or Mb. If steric constraints are of major importance, then  $\nu_{\text{O}_2}$  for HbZ and the others should approach that of the "picket fence" model; if polarity differences are responsible, then  $\nu_{\text{O}_2}$  of other Fe porphyrin complexes with altered pocket polarities would clarify the situation. The effect of different axial ligands on  $\nu_{\text{O}_2}$  would be of considerable interest as well.

Further work needs to be done on the effect of steric constraint on the axial base to delineate the difference in CO and O<sub>2</sub> binding properties of free porphyrin and base appended porphyrins. One such study is the CO, or preferably O<sub>2</sub>, binding to base appended porphyrins with different length attachments. Another approach is to use 2-MeIm or other hindered free bases, which cannot approach the porphyrin plane closely, in ferrous porphyrin ligand affinity studies. In fact, the excellent work of Rougee and Brault (38), in which Fe(deut)(2-MeIm) was found to bind CO some 200 times more poorly than Fe(deut)(Im), is the first piece of evidence that the Hoard-Perutz mechanism of cooperativity (62) is actually capable of explaining the decreased O<sub>2</sub> affinity of Hb in the tensed form.

## CONCLUSIONS

The nature of O<sub>2</sub> binding in Hb and Mb and in simple Fe porphyrin models is becoming clear. When O<sub>2</sub> is bound as a bent, rather than a triangular, ligand, it is best described as bound superoxide. This is true for Fe, Co, and perhaps as well for Cr. This bent geometry may be critical to biological functioning because it allows the discrimination between O<sub>2</sub> and CO. The use of steric bulk near the Fe binding site in hemoproteins to distort the FeCO geometry and hence lower its stability (without significantly affecting O<sub>2</sub> bonding) seems essential to avoid massive poisoning from the CO intrinsically formed from porphyrin catabolism.

$\ddagger$  The only other reported  $P_{1/2}^{\text{CO}}$  for free porphyrins are of appended base ferrous porphyrins (50, 51). The reported value of 0.2 torr has been revised (T. G. Traylor, personal communication) and is now in agreement with the results of ref. 38. This value is similar to that of microperoxidase (52) ( $P_{1/2}^{\text{CO}} = 0.0004 \text{ torr}$  at pH 7, 20°) a biological histidine-tailed ferrous porphyrin.

We thank Dr. Susan Hayes for synthesis of some of the compounds used, Peter Denisevich and Mark Nelson for stimulating discussions, the Fannie and John Hertz Foundation for fellowship support (K.S.S.), and the National Science Foundation Grant no. CHE75-17018, and the National Institutes of Health, Grant no. GM17880. Special thanks to Dr. Roy Gritter for his help.

- Collman, J. P., Gagne, R. R., Reed, C. A., Robinson, W. T. & Rodley, G. A. (1974) *Proc. Natl. Acad. Sci. USA* **71**, 1326-1329.
- Rodley, G. A. & Robinson, W. T. (1972) *Nature* **235**, 438-439.
- Collman, J. P., Gagne, R. R., Reed, C. A., Halbert, T. R., Lang, G. & Robinson, W. T. (1975) *J. Am. Chem. Soc.* **97**, 1427-1433.
- Tovrog, B. S., Kitko, D. J. & Drago, R. S. (1976) *J. Am. Chem. Soc.*, in press.
- Burness, J. H., Dillard, J. G. & Taylor, L. T. (1975) *J. Am. Chem. Soc.* **97**, 6080-6088.
- Hoffman, B. M., Szymanski, T. & Basolo, F. (1975) *J. Am. Chem. Soc.* **97**, 673-674.
- Goddard, W. A. & Olafson, B. D. (1975) *Proc. Natl. Acad. Sci. USA* **72**, 2335-2339.
- Loew, G. H. & Kirchner, R. F. (1975) *J. Am. Chem. Soc.*, **97**, 7388-7390.
- Dedieu, A., Rohmer, M. M., Bernard, M. & Veillard, A. (1976) *J. Am. Chem. Soc.* **98**, 3717-3718.
- Fantucci, P. & Valenti, V. (1976) *J. Am. Chem. Soc.* **98**, 3832-3838.
- Pauling, L. (1964) *Nature* **203**, 182-183.
- Weiss, J. J. (1964) *Nature* **202**, 83-84.
- Collman, J. P., Gagne, R. R., Gray, H. B. & Hare, J. W. (1974) *J. Am. Chem. Soc.* **96**, 6522-6524.
- Collman, J. P., Brauman, J. I. & Suslick, K. S. (1975) *J. Am. Chem. Soc.* **97**, 7185, footnote 4.
- Overend, J. (1963) in *Infrared Spectroscopy and Molecular Structure*, ed. Davies, M. (Elsevier Publishing Co., New York), pp. 345-376.
- Pinchas, S. with Laulicht, I. (1971) *Infrared Spectra of Labelled Compounds* (Academic Press, New York).
- G. Herzberg (1950) *Molecular Spectra and Molecular Structure. I. Spectra of Diatomic Molecules* (Van Nostrand Co., New York), p. 560.
- Maxwell, J. C., Volpe, J. A., Barlow, C. H. & Caughey, W. S. (1974) *Biochem. Biophys. Res. Commun.* **58**, 166-171.
- Barlow, C. H., Maxwell, J. C., Wallace, W. J. & Caughey, W. S. (1973) *Biochem. Biophys. Res. Commun.* **55**, 91-95.
- Maxwell, J. C. & Caughey, W. S. (1974) *Biochem. Biophys. Res. Commun.* **60**, 1309-1314.
- Crumbliss, A. L. & Basolo, F. (1970) *J. Am. Chem. Soc.* **92**, 55-60.
- Landels, J. D. & Rodley, G. A. (1972) *Synth. Inorg. Met.-Org. Chem.* **2**, 65-70.
- White, D. A., Solodar, A. J. & Baizer, M. M. (1972) *Inorg. Chem.* **11**, 2160-2167.
- Cheung, S. K., Grimes, C. J., Wong, J. & Reed, C. A. (1976) *J. Am. Chem. Soc.*, in press.
- Guilard, R., Fontesse, M., Fournari, P., Lecomte, C. & Protas, J. (1976) *J. Chem. Soc. Chem. Commun.* **1976**, 161-162.
- Valentine, J. S. (1973) *Chem. Rev.* **73**, 235-245.
- Siebert, H. (1966) *Schwingungs Spectroskopie in der Anorganischen Chemie* (Springer Verlag, Berlin), pp. 40, 51.
- Rolfe, R., Holzer, W., Murphy, W. F. & Bernstein, J. H. (1968) *J. Chem. Phys.* **49**, 963.
- Evans, J. C. (1969) *J. Chem. Soc. Chem. Commun.* **1969**, 682-683.
- Caughey, W. S. (1970) *Ann. N.Y. Acad. Sci.* **174**, 148-153.
- Antonini, E. & Brunori, M. (1971) *Hemoglobin and Myoglobin in their Reactions with Ligands* (American Elsevier, New York), pp. 224, 262, and 351.
- Roughton, F. J. W. (1954) *J. Physiol. (London)* **126**, 359-383.
- Huestis, W. H. & Raftery, M. A. (1975) *Biochemistry* **14**, 1886-1892.
- O'Toole, M. C. (1972) Ph.D. Dissertation, Arizona State University.
- Amiconi, G., Antonini, E., Brunori, M., Formanek, H. & Huber, R. (1972) *Eur. J. Biochem.* **31**, 52-58.
- Wallace, W. J., Volpe, J. A., Maxwell, J. C., Caughey, W. S. & Charache, S. (1976) *Biochem. Biophys. Res. Commun.* **68**, 1379-1386.
- Wang, J. H., Nakahara, A. & Fleischer, E. E. (1958) *J. Am. Chem. Soc.* **80**, 1109-1113.
- Rougee, M. & Brault, D. (1975) *Biochemistry* **14**, 4100-4106.
- Maxwell, J. C. & Caughey, W. S. (1976) *Biochemistry* **15**, 388-396.
- Brown, L. D. & Raymond, K. N. (1975) *Inorg. Chem.* **14**, 2595-2601.
- Gall, R. S., Rogers, J. F., Schaefer, W. P. & Christoph, G. G. (1976) *J. Am. Chem. Soc.*, in press.
- Barlow, C. H., Ohlsson, P. I. & Paul, K. G. (1976) *Biochemistry* **15**, 2225-2229.
- Lexa, D., Momenteau, M. & Mispelter, J. (1974) *Biochim. Biophys. Acta* **338**, 153-163.
- Huber, R., Epp, O. & Formanek, H. (1970) *J. Mol. Biol.* **52**, 349-354.
- Padlan, E. A. & Love, W. E. (1975) *J. Biol. Chem.* **249**, 4067-4078.
- Norvell, J. C., Nunes, A. C. & Schoenborn, B. P. (1975) *Science* **190**, 568-570.
- Hoard, J. L. (1975) *Porphyryns and Metalloporphyryns*, ed. Smith, K. M. (Elsevier Scientific Publishing Co., New York), pp. 356-358.
- Caughey, W. S., Alben, J. O., McCoy, S., Boyer, S. H., Charache, S. & Hathaway, P. (1969) *Biochemistry* **8**, 59-62.
- Morimoto, H., Lehman, H. & Perutz, M. (1971) *Nature* **232**, 408-413.
- Chang, C. K. & Traylor, T. G. (1973) *J. Am. Chem. Soc.* **95**, 8475-8477.
- Geibel, J., Chang, C. K. & Traylor, T. G. (1975) *J. Am. Chem. Soc.* **97**, 5924-5926.
- Sharma, V. S., Ranney, H. M., Geibel, J. F. & Traylor, T. G. (1975) *Biochem. Biophys. Res. Commun.* **66**, 1301-1306.
- Caughey, W. S. (1970) *Ann. N.Y. Acad. Sci.* **174**, 148-153.
- Robinson, E. & Robbins, R. C. (1970) *Ann. N.Y. Acad. Sci.* **174**, 89-95.
- Sjostrand, T. (1952) *Acta Physiol. Scand.* **26**, 328-344.
- Metz, G. & Sjostrand, T. (1954) *Acta Physiol. Scand.* **31**, 384-342.
- O'Earra, P. (1975) *Porphyryns and Metalloporphyryns*, ed. Smith, K. M. (Elsevier Scientific Publishing Co., New York), pp. 123-153.
- Lemberg, R. (1935) *Biochem. J.* **29**, 1322-1338.
- Coburn, R. F. (1970) *Ann. N.Y. Acad. Sci.* **174**, 11-22.
- Gibson, Q. H., Heller, P. & Yakulis, V. (1966) *J. Biol. Chem.* **241**, 1650-1651.
- Winterhalter, K. H., Anderson, N. M., Amiconi, G., Antonini, E. & Brunori, M. (1969) *Eur. J. Biochem.* **11**, 435-440.
- Perutz, M. F. (1970) *Nature* **228**, 726-739.